

ACCESSION NR: AP4040487

Elementoorganic Compounds, AN SSSR)

SUBMITTED: 11Jul63

DATE ACQ: 06Jul64

ENCL: 00

SUB CODE: GC

NO REF SOV: 003

OTHER: 008

Card 3/3

ACCESSION NR: AP4040488

S/0190/64/006/006/1087/1091


AUTHORS: Korshak, V. V.; Krongauz, Ye. S.; Berlin, A. M.; Travnikova, A. P.

TITLE: Synthesis of polymers by the polycyclization reaction. 6. Polypyrazoles

SOURCE: Vysokomolekulyarnyye soyedineniya, v. 6, no. 6, 1964, 1087-1091


TOPIC TAGS: polycyclization reaction, polypyrazole, bipyrazole polycondensation, dicarboxylic acid chloride, diketone polycyclization, dicarboxylic acid dihydrazide

ABSTRACT: The investigators attempted to synthesize polypyrazoles from compounds containing pyrazole cycles. The desired results were achieved by polycondensation of bipyrazoles with the chlorides of dicarboxylic acids according to the reaction



Card 1/3

ACCESSION NR: AP4040468



where X = C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>; C<sub>6</sub>H<sub>4</sub>OC<sub>6</sub>H<sub>4</sub>; CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>; (CH<sub>2</sub>)<sub>8</sub>; R = CH<sub>3</sub>, C<sub>6</sub>H<sub>5</sub>;  
Y = (CH<sub>2</sub>)<sub>4</sub>, C<sub>6</sub>H<sub>4</sub>.

A total of 8 bypyrazoles were synthesized. Seven of them were new and represented: 4,4'-bis-(5-methylpyrazolyl-3)diphenyloxide, 4,4'-bis-(3,5-dimethylpyrazolyl-4)xylilene, 4,4'-bis-[(3,5-dimethylpyrazolyl-4)methyl]diphenyloxide, 4,4'-bis-[(3,5-dimethylpyrazolyl-4)methyl]diphenyl, 1,8 di-(5-phenyl-pyrazolyl-3)octane, di-(3,5-dimethylpyrazolyl-4), and 4,4'-bis-(5-methylpyrazolyl-3)diphenyldisulfide. The procedure was started by mixing 30-40 ml of pyridine with 0.1 mole quantities of one of the bypyrazoles. To these mixtures were added (dropwise) 0.1 mole amounts of adipic, terephthalic, or isophthalic acid chloride, dissolved in 20 ml of xylene. The contents of the flasks were stirred and cooled for several hours. They were then heated for a long time to 100-125°C, and were allowed to stand overnight. The polypyrazoles so produced were identical with the polypyrazoles ob-

Card 2/3

ACCESSION NR: AP4040488

tained by polycyclization of bis-( $\beta$ -diketones) with the dihydrazides of the corresponding dicarboxylic acids. The latter group was described in an earlier publication by the authors and P. N. Gritkova (Dokl. AN SSSR, 148, 602, 1963). Orig. art. has: 3 tables and 1 formula.

ASSOCIATION: Institut elementoorganicheskikh soedineniy AN SSSR (Institute of Elementoorganic Compounds, AN SSSR)

SUBMITTED: 11Jul63

DATE ACQ: 06Jul64

ENCL: 00

SUB CODE: GC

NO REF SOV: 004

OTHER: 006

Cord 3/3

ACCESSION NR: AP4042185

S/0190/64/006/007/1195/1202

AUTHOR: Korshak, V. V.; Krongauz, Ye. S.; Berlin, A. M.; Smirnova, T. Ya.

TITLE: Synthesis of polymers by polycyclization. Polypyrazoles. VII.

SOURCE: Vy\*sokomolekulyarny\*ye sovedinaniya, v. 6, no. 7, 1964, 1195-1202

TOPIC TAGS: polypyrazole, polycyclization reaction, bis-( $\beta$ -diketone), dihydrazine, hexamethylenhydrazine dihydrochloride, p-phenylenehydrazine dihydrochloride, polypyrazole property

ABSTRACT: The authors have synthesized polypyrazoles (mp, . 200—300C) by polycyclization of linear and branched bis-( $\beta$ -diketones) with dihydrazides of dicarboxylic acids. In an attempt to develop polypyrazoles with a higher heat resistance, dihydrazides were replaced with dihydrazine, or amide groups were introduced in the polymers to form hydrogen bonds. Polycyclization of bis-( $\beta$ -diketones) with hexamethylene- or p-phenylenhydrazine dihydrochlorides in boiling alcohol with alkali added to separate and bind HCl, or heating equimolar amounts of the initial materials in pyridine, yielded

Cord 1/2

ACCESSION NR: AP4042185

polypyrazoles — powders with a mp of 80—265C and a molecular weight of 5000. Polypyrazoles containing amide groups in the backbone were synthesized by reacting dipyrazoles with diisocyanates in chlorobenzene or by melting the initial materials in nitrogen. These polymers are white powders with a mp of 208—276C and a molecular weight of up to 10,000. IR spectra indicate that they do not contain hydrogen bonds. Thus, the attempt to synthesize heat-resistant polypyrazoles failed. The presence of heavy pyrazole rings upsets the symmetry and loosens the packing density of the polymer chains, and, as a result, prevents the formation of hydrogen bonds. Orig. art. has: 1 figure and 2 tables.

ASSOCIATION: Institut elementoorganicheskikh soedineniy AN SSSR  
(Institute of Organoelemental Compounds, AN SSSR)

SUBMITTED: 11Jul63

ATD PRESS: 3068

ENCL: 00

SUB CODE: OC, GC

NO REF SOV: 009

OTHER: 003

Card 2/2

REEL #49

BEREZOVA, ye. F.  
TO

BERLIN, A.M

END